

IOWA STATE UNIVERSITY

Digital Repository

Integrated Crop Management News

Agriculture and Natural Resources

9-13-2013

Iowa Farmer Perception of Corn Rootworm Resistance

Erin W. Hodgson

Iowa State University, ewh@iastate.edu

Kristine J. P. Schaefer

Iowa State University, schaefer@iastate.edu

Aaron J. Gassmann

Iowa State University, aaronjg@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Entomology Commons](#)

Recommended Citation

Hodgson, Erin W.; Schaefer, Kristine J. P.; and Gassmann, Aaron J., "Iowa Farmer Perception of Corn Rootworm Resistance" (2013). *Integrated Crop Management News*. Paper 19.
<http://lib.dr.iastate.edu/cropnews/19>

This Article is brought to you for free and open access by the Agriculture and Natural Resources at Digital Repository @ Iowa State University. It has been accepted for inclusion in Integrated Crop Management News by an authorized administrator of Digital Repository @ Iowa State University. For more information, please contact digirep@iastate.edu.

[ICM Home](#)[ISU Extension Calendar](#)[Publications](#)[Extension News](#)[County Offices](#)[Contact Us](#)

search

Subscribe to Crop
News

Archives

[2014](#)[2013](#)[2012](#)[2011](#)[2010](#)[2009](#)[2008](#)[Previous Years](#)

ISU Crop Resources

[Extension Field
Agronomists](#)[Crop & Soils Info](#)[Pesticide Applicator
Training](#)[Agronomy Extension](#)[Entomology Extension](#)[Plant Pathology
Extension](#)[Ag and Biosystems
Engineering Extension](#)[Agribusiness
Education Program](#)[Iowa Grain Quality
Initiative](#)[College of Agriculture
and Life Sciences](#)[ISU Extension](#)

Integrated Crop Management NEWS

PRINT STORY
EMAIL STORY
ADD TO DELICIOUS
ATOM FEED
FOLLOW ON TWITTER

Iowa Farmer Perception of Corn Rootworm Resistance

By Erin Hodgson, Kristine Schaefer and Aaron Gassmann, Department of Entomology

Every year Iowa State University (ISU) Extension and Outreach provides private pesticide applicator training covering a variety of pesticide safety and pest management issues. In response to reported incidences of Bt-resistant corn rootworm, a series of remote "clicker" questions were developed to survey current corn rootworm management practices and assess the degree to which farmers perceived resistance in 2012. The questions were asked at 153 pesticide applicator training sessions facilitated by extension field agronomists. More than 8,100 meeting participants responded to the six multiple-choice questions. Responses were anonymous and not everyone answered every question. Here are the statewide summaries of selected questions from the survey based on percentages of respondents:

To ascertain current management practices, participants were asked how they manage corn rootworm (Figure 1). Of the various management options for corn rootworm, 17 percent of respondents used Bt traits, 25 percent used crop rotation and 38 percent responded they used three or more methods.




Figure 1. How do you manage corn rootworm?

When asked if they had fields in 2012 where they suspected Bt trait failure, almost one-fourth (23 percent) of participants responded "yes" and 43 percent responded that they reported the suspected failure to someone. The majority of respondents who suspected trait failure either dug and rated root injury (38 percent) or noted goosenecking/lodging symptoms (33 percent) to assess the injury (Figure 2). About 13 percent of people who suspected a Bt trait failure were able to verify the larval injury (Figure 3). Although lodging can occur as a result of rootworm feeding injury, there may be other causes such as strong wind events. The only way to verify Bt trait performance issues is to evaluate root injury. Thus the rate of verified trait failures from this survey may be overestimated because larval root injury was not confirmed for all cases.

Additionally, the level of root injury that a grower may have considered sufficiently high to warrant a trait failure is unknown. Generally, a trait failure would be considered to be present if there is one node of root injury for Bt corn with a single trait targeting rootworm and 0.5 nodes for corn that has two Bt traits targeting rootworm. A [root rating scale](#) developed by the Iowa State University Department of Entomology is available.




Figure 2. What method did you use to assess corn rootworm injury?




Figure 3. What was the outcome of the assessment?

To understand where most of the perceived Bt trait failures were occurring within Iowa, we grouped responses by region. About one-third of people in northeast Iowa suspected a Bt trait failure in 2012 (Figure 4). Most of the positive verifications were in north central Iowa, with approximately one-third of people reporting Bt trait failures (Figure 5).




Figure 4. Percent of respondents who suspected a corn rootworm trait failure in at least one of their cornfields in 2012.




Figure 5. Percent of respondents who positively verified corn rootworm trait failure.

If you find injury to Bt corn on more than one node, consider diversifying pest management for that field. Severe injury may be due to western corn rootworm that has developed Bt resistance, and resistance should be suspected in these cases. The Department of Entomology at Iowa State University is working to understand the extent of resistance within the state and to develop management recommendations for Bt-resistant populations of western corn rootworm. Please contact Aaron Gassmann or Erin Hodgson to report fields with suspected resistance.

Erin Hodgson is an associate professor of entomology with extension and research responsibilities; contact her at ewh@iastate.edu or 515-294-2847. Kristine Schaefer is program manager of the pest management and the environment program and can be reached at schaefer@iastate.edu or 515-294-4286. Aaron Gassmann is an assistant professor of entomology with research and teaching responsibilities in insect pest management; contact him at aaronjq@iastate.edu or 515-294-7623.

This article was published originally on 9/13/2013. The information contained within the article may or may not be up to date depending on when you are accessing the information.

Links to this material are strongly encouraged. This article may be republished without further permission if it is published as written and includes credit to the author, Integrated Crop Management News and Iowa State University Extension. Prior permission from the author is required if this article is republished in any other manner.

Copyright ©2014 [Iowa State University Extension](#) | [Iowa State University](#)
[Contact us](#) | [For Staff](#) | [Nondiscrimination and Information Disclosures](#) | [CMS Admin](#)
Last Updated 9/14/2013